

LESPEDEZA WEED CONTROL

Lisa Baxter, Extension Forage Specialist

HERBICIDE	BROADCAST RATE/ACRE		REMARKS AND PRECAUTIONS
	AMOUNT OF FORMULATION	POUNDS ACTIVE INGREDIENT	
<b>PREPLANT INCORPORATED</b>			
EPTC Eptam 7 lb/gal	3.5 pt	3.1	For winter annual grass and some broadleaf weed control, incorporate 2–3" deep before planting for control of annual grass and broadleaf weeds. Do not use if a grass or grain crop is to be planted with the lespedeza.
<b>POSTEMERGENCE—DORMANT APPLICATIONS</b>			
Gramoxone SL 2.0 others	1–2 pt	0.25–0.5	Apply during winter months when lespedeza is dormant. Controls Italian ryegrass, common chickweed, and most other winter annual broadleaf weeds. Applications to lespedeza that is not dormant may result in stand and/or yield reductions. Add a nonionic surfactant at 1 pt/100 gal of spray solution. <b>DO NOT</b> graze or cut for hay within 60 days of application. Make only one application per season.
2,4-Dichlorophenoxyacetic acid + florpyrauxifen-benzyl  NovaGraz	24–48 oz	0.50+0.02 — 1.00+0.01	Provides good control of multiple broadleaf weeds. Add a methylated seed oil at 1 gal/100 gal of spray solution. Annual lespedeza will exhibit some initial injury (such as lodging and loss of vigor) but will recover. Do not cut or harvest grass for forage or hay within 14 days after application. Do not graze dairy cattle on treated areas for 3 days after application. Do not graze meat animals on treated areas within 3 days before slaughter. Do not make more than two applications per year.